

JULIE FAST: Behind the mood-swing mask

HOPE AND HARMONY FOR PEOPLE WITH BIPOLAR



**THE
POWER
OF
TWO**
**TACKLING
BIPOLAR WITH
YOUR PARTNER**



**HOPE
HEALS**

DEPRESSION
in mind
&
body

10 strategies
for
**SMOOTHER
SAILING**

**Clutter
OUT →
Harmony
IN ←**

the power of two

Your partner can become a **pillar of strength** when you **work together as a couple**

Written by Barbara Boughton

There's no doubt that the nature of your relationship with a "significant other" has a major effect on managing bipolar. Research now shows that having a supportive partner may be just as important as medication and psychotherapy in preventing relapse, according to Sagar Parikh, MD, a professor of psychiatry at the University of Toronto.

Likewise, the stress of high-conflict relationships or the emotions that accompany breakup and divorce can sometimes exacerbate symptoms of the disorder, Parikh says.

So what can you do to make your partner an ally in recovery?

The first step, says Parikh, is education—for both of you.

The more that both partners know about symptoms, treatments, and coping strategies, the more they can work together to address common challenges.

Reading and online research, workshops presented by mental health organizations, discussions with mental health practitioners, and peer support groups are all good ways to get informed.

A partner or spouse who is up to speed on what it takes to live with bipolar will find it easier to understand when you ask for support.

The next step is learning to discuss matters relating to your illness openly and honestly.

For one thing, being able to share what's going on in your life and your head provides your partner with a context for any irritability, sadness or high spirits you exhibit. For another, it gives you both a touchstone for recognizing early signs of a mood shift.

Elizabeth Barr and her husband, Rory, (*pictured at left*), who have been married since August 2012, have a conversation at least once a week about any symptoms Elizabeth might be experiencing.

Learning when a hug can help

When **Heather Tobin-Lambert and Ken Lambert** of Merrimack, New Hampshire, first met in fall 2002, Heather certainly wasn't in the mood for romance. She was experiencing serious depression and "very resistant to dating," but Ken waited patiently.

"When we did start dating, I was very forthright about having bipolar disorder, and he was very understanding," recalls Heather, 44.

Now Heather's relationship with her husband is a rock that steadies her when symptoms crop up. In Heather's case, that's often behaviors that signal depression—not getting out of bed, failing to eat healthy foods, and crying a lot.

After nine years of marriage, Ken has come to understand what his wife needs when she is ill. And Heather made it a priority to communicate to her husband about what kind of support she finds the most helpful.

"Sometimes it's just the simple things that are important," Heather says. "A hug from Ken can be very grounding when I'm having a difficult time. He also knows when to give me space, such as when I get irritable, angry, and when I lash out."

Perhaps more importantly, Ken and Heather have an agreement that they'll promptly address any worrisome symptoms.

Heather admits she's not always receptive when Ken spots signs of depression or hypomania, such as being quick to anger or spending too much on online shopping, yet she will usually agree to call her doctor.

"Couples therapy has also helped us," she says. "We've learned not to blame each other when we discuss sensitive issues such as my bipolar disorder, and to respectively express our feelings. Sometimes that may mean taking a break or even going to different rooms when a discussion becomes too heated."

She adds, "What has been most important is for both of us to realize that I'm more than the identified patient in our relationship. I'm Heather, a wife who is more than her illness, and who needs and wants hope and love from her partner."




Photo: DACHOWSKI PHOTOGRAPHY.COM

"Regular communication is really important," says Elizabeth, 32, of New Westminster, British Columbia. "We talk about what I'm feeling and things that he notices about me. Sometimes, it's hard for me to see the forest for the trees, especially if I'm not feeling well."

Rory's feedback provides her with a reality check, Elizabeth says.

"Last year I had a manic episode and Rory realized something was wrong when I told him: 'I want to go on a 5K run.' I'm a pretty sedentary person, so for me that's out of character. It gave Rory a clue that I might be experiencing mania," she recalls.

In most intimate relationships, it's important to make significant others aware of red flags, according to David Miklowitz, PhD, a professor of psychiatry and direc-

tor of the Child and Adolescent Mood Disorders Program at the Semel Institute for Neuroscience and Human Behavior at the University of California—Los Angeles.

“People with bipolar disorder can make a list of symptoms and behaviors that they know indicate early signs of a manic or depressive episode. The partners can then refer back to these lists in order to spot early symptoms,” Miklowitz says.

your medicines? Some people like that kind of active support from their partners, and some do not.”

In some cases, Miklowitz says, intervention from a spouse can be seen as “too parental or too controlling.”

The bottom line is this: In order to tell your partner how to be helpful, you first need to know what kind of help you want. That’s highly individual, so every couple

and our relationship. She knows that I can stand on my own two feet, and she doesn’t judge me,” says Bill, a retired refrigeration pipe fitter.

What Bill appreciates are Telle’s understanding and compassion for his symptoms, her trust that he is working to get the best treatment he can, and her emotional support when he’s had to be hospitalized.

Still, Telle has occasionally taken an

“Regular communication is really important. We talk about what I’m feeling and things that he notices about me.”

comfort level

With education and experience, partners often become adept at spotting subtle signs, Miklowitz notes—sometimes before the person with bipolar does.

“Then the question becomes: If your partner spots early signs of mania or depression, what do you want them to do with that information?” Miklowitz says.

“Do you want them to call your doctor? Do you want them to go in with you for a therapy session, or encourage you to take

will find a different fit. After a disturbing interaction with a friend or boss, one person may simply want to vent while another seeks advice on how to resolve the situation.

Bill Scholtz, 67, has never looked for a lot of hands-on involvement from his wife, Telle. The couple lives near Santa Cruz, California, and has been married for 20 years.

“The main way that Telle has supported me is through her acceptance of me—and that’s been very important to me

active role in Bill’s care. In May 2012, Bill became delusional while being assessed at a hospital for a blood clot in his lung. Telle spoke to his psychiatrist, who arranged for Bill to be admitted for psychiatric intervention.

If your partner does take an active role, Miklowitz says, it’s important to understand that the person will probably make mistakes.

“It’s crucial to give your partner some leeway, especially in the beginning of a relationship, when they’re just getting to

know you and the effects of your illness. They're not trying to control you. They're just trying to help, and they don't always know how to best do that," he says.

talking points

Once you are clear in your own mind about what role you'd like your partner to take, Miklowitz notes, it's crucial to be direct in

Mental Health in Montreal, stresses the need to express your wants and feelings with compassion.

With mirroring, for example, first reflect your partner's viewpoint, then express understanding and empathy to validate how the other person feels.

In addition, she says, "both partners should be careful to use 'I statements' rather than blaming the other person in talking about the bipolar disorder. The im-

As a further step, Poirier notes, you and your partner can write a contract about what each of you will do when warning signs appear. A document you signed when well may be more effective against denial when you're ill than a partner's reminders.

There are other small but helpful ways partners can counter symptoms, she says. During a manic phase, reduce stimulation by decreasing light and loud music in the household, and keep credit and cash cards

Then the question becomes: If your partner spots early signs of mania or depression, what do you want them to do with that information?

expressing your wishes—and also to listen to what your partner has to say.

Miklowitz recommends asking "clarifying questions" to make sure you understand your partner's point of view. To doublecheck that you're interpreting correctly, practice repeating back what the other person said—a technique known as reflective listening or mirroring.

Nancy Poirier, MPs, a clinical psychologist at the Douglas University Institute for

portant thing is for both partners to feel understood."

As far as practical approaches, Poirier recommends developing a relapse prevention plan that details early warning signs of a mood shift and how to respond. The Barrs' plan includes a list of people whom Rory should call to get Elizabeth treatment and/or medication, as well as the name and contact information for the hospital where she'd like to be treated.

inaccessible. During a depressive phase, let more light into the house, encourage exercise, and gently urge completion of at least one pleasant activity each day.

On a day-to-day basis, agree on a system that keeps encouragement from feeling like nagging. Again, this will vary from couple to couple. Some might be comfortable with a daily check-in about medication compliance, for example, while others will prefer a more emotionally neutral chart



Photo: Courtesy of DAN & DOLORES SEGAL

In sickness and in health

Dan and Dolores Segal of St. Louis, Missouri, have weathered 43 years of marriage—and Dan’s bipolar swings.

Dan, now 73, was accurately diagnosed in his late 20s. He and Dolores were contemplating marriage, and she accepted the news along with the man.

“He was a real charmer and had a wonderful sense of humor,” Dolores remembers.

Dan was in a stable period when they met, but shortly after the wedding he slipped into depression. For the next decade, he cycled from months of mania marked by reckless spending and anger, to months of immobilizing depression, to stretches of stability that might last half a year. He went through a series of jobs, never keeping one for long.

“I was ill and non-productive and she was the provider and caretaker,” Dan says.

Dolores held down several part-time jobs, taking their two children to work with her because they couldn’t afford child care. The couple went through two bankruptcies.

“It was painful and it was devastating, but I didn’t leave because I don’t believe in divorce,” says Dolores, 75. “But even in the most difficult moments, Dan’s sense of humor would come shining through.”

They held on until Dan found the treatment he needed at Washington University. An equally important turning point was their decision to start a peer support group in 1980, which became one of the founding chapters of what is now the national Depression and Bipolar Support Alliance (DBSA).

The work not only helped his wife become more educated about the disorder, Dan says, but through running meetings “she gained many skills in talking with me in a very helpful and therapeutic way.”

“We learned how to communicate even around tough topics—and not blame the other person,” Dolores adds.

Dolores admits to “mixed feelings” about their marriage over the years, but says that their life improved as their ability to address Dan’s health got better.

“He has changed in a really positive way over the years,” she says. “And our communication strategies just got better and better. We feel so blessed now.”

system that both partners can refer to.

Couple’s therapy can be an effective way to develop strategies for coping with the disorder together, Miklowitz says.

“The counseling should help the couple solve problems that arise around the symptoms of bipolar disorder, and learn to communicate effectively about them,” he says.

In addressing specific issues, Miklowitz adds, it helps to realize that you may not get everything you want.

“You have to be willing to compromise,” he says. “You have to start with the assumption that everyone—you and your partner—are doing their best.”

“Doing your best” might involve taking as much responsibility as you can for your recovery and well-being, Poirier points out. For your partner, that might mean cutting you a little slack for symptomatic behavior.

Elizabeth Barr says that when she succumbs to what she calls her “ooh, shiny!” syndrome, Rory usually accepts the impulsive action as part of her illness.

When she bought several hundred dollars’ worth of tea towels, she recalls, Rory

Another definition of partner

People who are single may find a family member who becomes a partner in recovery. For **Jackie Malmstrom** of Evans City, Pennsylvania, that's her dad, **Larry Nulton**.

"We've become an amazing team at managing my bipolar I," explains the 25-year-old. "We even have a special binder that we keep on the coffee table that has copies of the most important papers about my illness and how we should handle things."

The binder includes a list of her medications and what they're for, contact information for her doctors, steps to take in various crisis situations, even copies of some of her homework from psychotherapy.

Malmstrom has found the skills she's learning in dialectical behavior therapy especially helpful in managing her symptoms. When she comes home from group, she shares new information with her father. Also really helpful, she says, is "both of us reading everything we can get our hands on about bipolar. We test out new techniques for managing things to see what works best for us."

For example, Malmstrom now keeps track of when (or whether) she's taken her medications on a dry-erase board that hangs on the fridge. The chart not only reminds her to take the meds, but also keeps her father in the loop without having to bug her.

Malmstrom, who was diagnosed at 14, has seen a huge change since her father started educating himself. When she's in a mood shift, they're able to talk through what may have triggered it and how to get her back in balance. When her dad learned that people with mental illnesses sometimes use cigarettes to cope with stress, he stopped insisting she quit smoking. Instead, he encourages her to cut back.

"Before he started reading he didn't understand nearly as well as he does now, which really goes without saying," she reflects. "We fought a lot more and he really didn't 'get' me. Now we fight less and he's much more compassionate to what I'm enduring."




Photo: ALWAYS REMEMBER PHOTOGRAPHY/MARSHA LONGCHENNA

commented that they were now stocked up for a few years. When she took their cat to the vet and came home with an extra kitten, Rory "accepted the kitten with open arms," she reports—and they now have a rule that they only go to the vet's together.

"He isn't a pushover," she says, "he just knows that in the spring, I pull a few whoppers and that is OK."

In fact, she says, Rory took the news of her diagnosis better than she did. They were dating at the time, Elizabeth says, and Rory made it a point to find books that would be helpful.

"He never judged me for having bipolar," she adds. "I'm very lucky to have found Rory." ●

Barbara Boughton, a freelance health and medical writer, writes for Reader's Digest, Medscape.com and other outlets. She lives in the San Francisco Bay area.